

**Effect of the entomofungns *Beauveria bassiana* on different stages of Angoumois moth *Sitotroga cerealella***Hussam Aldin Abdullah Mohammad , Safaa Zakaria Baker , Hind Numan Harhosh  
Plant Prot., Agric. College, Baghdad Univ., Iraq. E-mail: Mobil No.: 07707797789

## معلومات البحث

تاريخ استلام البحث  
2016/7/7  
تاريخ قبول البحث  
2016/9/7

## Keywords

Beauveria  
bassiana ،  
Sitotroga  
cerealella

## Abstract

This experiment was conducted in 2016, to investigate the effect of entomofungns *Beauveria bassiana* on different stages of Angoumois moth *Sitotroga cerealella*. The Results showed that eggs hatching percentage was zero % in three concentrations of fungus *Beauveria bassiana*, while in the control treatment was 96%. The average of larval mortality was 16.0 % in the concentration  $1 \times 10^6$  spore/ml and the lowest percentage Larval mortality was 3.0 % in control treatment. The lowest a verge of pupa emergence was 8.3% in concentration  $1 \times 10^6$  spore /ml, while heighest in control was 96.3 %.The height mortality of adult percentage in the concentration  $1 \times 10^6$  spore /ml was 96.6 % , while in control treatment was 16.6 % after 96 hours at the fungus treatment.

**تأثير الفطر الممرض *Beauveria bassiana* على الاطوار المختلفة لعتة الانجومو *Sitotroga cerealella***

حسام الدين عبد الله محمد ، صفاء زكريا بكر ، هند نعمان حرحوش

## المستخلص

في تجربة اجريت عام 2016 في مختبر الدراسات العليا / كلية الزراعة - جامعة بغداد - اوضحت النتائج ان نسبة فقس البيض المعامل بمعلق الفطر *B.bassiana* وبالتركيز الثلاثة بلغت صفر % بينما في معاملة المقارنة بلغت 96% ومعدل القتل لليرقات 16.0% في التركيز  $106 \times 1$  بوغ / ملتر اما اقل نسبة قتل فكانت في معاملة المقارنة والتي بلغت 3.0%. اقل معدل لخروج البالغات من العذارى المعاملة فكانت 8.3% بالتركيز  $106 \times 1$  بوغ / ملتر اما اعلى معدل لخروج للعذارى بلغت 96.3% بمعاملة المقارنة . اعلى معدل للقتل بالغات فان بالتركيز  $106 \times 1$  بوغ / ملتر فان 96.6% اما اقل معدل للقتل فان 16.6% في معاملة المقارنة بعد 96 ساعة في معاملة الفطر

\*Corresponding author : E-mail [salihhussamaldin@gmail.com](mailto:salihhussamaldin@gmail.com)

Al- Muthanna University All rights reserved.

## Introduction

The wheat plant *Triticum aestivum* is considered an important crops, its used in human food as bread,biscuits, pastures and many foods spread in the world(Hudson *et al.* 1981), the wheat consider meain food resource of human and domestic animals it has heightcalorie of carbohydrate and provide for adults more than 50 % from needs of proteins (Al- younis 1993).

Wheat was attacted by many pest caused damage to 55 % from main products in some areas (fornal *et al* 2007) . from these pests caused damage are the insects and important once was Angoumois

moth *Sitotroga cereallels* it's consider an important insects and widely spread in the world and caused damage to stored seeds (Togola *et al.* 2010) .

The grain moth attacked the plant intitle field when it's matntre stage and transferred with the infected wheat seeds to stores, itsconsider primary insects and its attack mainly wheat , barley , Rice , corn , sweet corn , and another stored crops (Howlaidr, Matin 2014) . The grain moth belong to order Lepidoptera, the Larvae feed on cereals that consume from and human its

causes losses in the crops about 40% from all stored cereals (Bushra and Astam 2017) .

We know that pesticides caused more damage to environment and non-target organism so the scientists found promising alternative pesticides that less harm to human and friendly ecology ( Kaur *et al.* 2011) . The most promising methods used the entomofungus like *Beauveria bassiana* that attack about 100 species from insects, and easy to producing and culturing on many artificial media and forming many conidia in million numbers which infect insects causing diseases called white muscarden so suggestion this fungus to control angoumois moth in cereal stored. Biological control the use of living organisms parasitoid , predators , pathogen and competitors to suppress pest population below levels that would occur naturally . Biological control programs are intended to suppress invasive through use the entomofungus like *b.bassiana* to control economic insects. This fungus has been attacked about 700 insects in culding Angoumois grain *S.cerealella*

#### **Materials and Methods:**

This experiments was Conducted in higher studies laboratory of Entomology in college of agriculture, Baghdad University in the year 2016.

The culture of Angoumois moth was carried from ministry of Science and technology, while the fungus is product from IPM program ministry of agriculture Iraq.

The Culture placed in incubator at  $28 \pm 2$  C° and R.H  $70 \pm 5$  % in all experiment below.

The percentage of fungus spores was  $1 \times 10^7$  spore /ml , one gram of powdered fungus dissolved in

one liter of distilled water stirrer for 10 minutes and leaved to one hour . pull up one milliliter of fungus suspension and added to test tube contains 9 ml distiller water ,then the concentration was  $1 \times 10^6$  spore /ml then pull up from this suspension one ml and added to test tube contain 9 ml distilled water then the concentration was  $1 \times 10^5$  spore / ml then we get a new concentration  $1 \times 10^2$  spore / ml (Lacey 1997).

#### **Effect of suspension of *B.bassiana* on different stags of Angoumois moth *S.cerealella***

Ten eggs were used 30 eggs as treatment and with three replicates for each treatment. three and control treatment with water only. Spray the eggs with three concentrations  $1 \times 10^2$  and  $1 \times 10^4$  and  $1 \times 10^6$  spore /ml using hand sprayer capacity 100ml from distance 15 cm . then the eggs that put in petri dish (9 cm) were treated by fungus suspension treated eggs placed in petri dishes 9 cm diameter, recorded percentage of mortality of eggs and followed still adult emergence. In the same way treated the first larval instar of Angoumois moth *S. cerelella* with fungus suspension in the same conc. And replication then put the larvae in Petri dish 9 cm and put 5 gram of grindery wheat for feed larvae recording the number of larva mortality daily for 30 days. Then Put in incubators in  $28 \pm 2$  C° and  $70 \pm 5$  % humidity.

#### **Treatment of pupae**

Five pupae of *S.cerelella* with three replicate for each treatment, the pupa spray 10 ml of suspension of fungus *B. bassiana* in the three concentrations  $1 \times 10^2$  ,  $1 \times 10^4$  and  $1 \times 10^6$  spore /ml while the control spray with water only . the

number of dead and alive pupae recorded until became adults .

**The adult's treatment**

Five paris of adult (male and female) were taken in the age 24 hours after emergence from pupa sprays in the three conc. of fungus suspension taken 10 ml in hand sprayer the contral spray with water only . every Petri dish added 5 gram of grindery wheat to make adults laying eggs on the cereal. the number of dead and alive of adults recorded for 10 days .

**Statically analysis: -**

Data were analyzed using SAS program with CRD experiment compered by L.S.D at 0.05 Level ( SAS 2012)

**Results and Discursions: -**

Table (1) refes that percentage of eggs hatching in three concentration of fungus was zero % while in controled was 96.0 % the fugus concentration caused mortality to eggs be that shell of eggs is not contain any antifungal material it caused essay

penetration the conidia to in egg and the conidia can also penetrate from micropyle .

Navon (2000) stated the shell of insects eggs consisted of many proteins and saccharide and some eggs have one micropyle like diptera insects and another have 35-45 micropyle like locust .

Vincent (2009) found eggs shell consisted of vitelline and its suitable for germination for conida of *B.bassiana*.

Galín *et al.* (2009) mentiaed the effect of conidia fungus *Metarhizium anisopliae* on hatching percentage due to present polar and not polar Lipids in the exolayer of egg shell that cause to germinate conidia of fungus because this material resource of energy to forming the appressorium how make the conidia stable un cuticle then penetrate to egg and feed on egg contain.

Mohamd (2007) reported the conidia of fungus *B.bassiana* caused 100 % mortality to eggs of mosquitoes *culex pipiens* in all concentration used.

Table (1). Effect of suspension of *B.bassiana* on hatching

Treatment	Hatching% of eggs		
	Exposure time(day)		
	2	4	6
Control	10.0	70.0	16.0
1x10 <sup>2</sup>	0.0	0.0	0.0
1x10 <sup>4</sup>	0.0	0.0	0.0
1x10 <sup>16</sup>	0.0	0.0	0.0
LSD 0.05	9.41	9.41	5.43

**Effect of suspension of fungus *B.bassiana* on first larval instar of angoumious moth *S.cerealella***

Table (2) show mortality of first instar on grindery wheat when treated in concentration  $1 \times 10^2$ ,  $1 \times 10^4$ ,  $1 \times 10^6$ , spore /ml the heightpercentage of mortality with concentration  $1 \times 10^6$  spore /ml was gave 16.8 % while in concentration  $1 \times 10^4$  spore /ml was gaved 9.6% in concentration  $1 \times 10^2$  spore /ml was 4.0 %, the statically found highl significantbetween concentrations and control .

The percentage of distorted Larvae was height in concentration  $1 \times 10^6$  spore /ml it was 72.2 % while in concentration  $1 \times 10^4$  spore /ml was 52.2 % the percentage of distorted lavae in control was zero.

The percentage of pupa was in control 69.6% while in concentration  $1 \times 10^6$  spore /ml was zero % .

The adult , emergence was 48.0 % in control treatment while in  $1 \times 10^6$  and  $1 \times 10^4$  spore /ml was

zero % . That may be due to that first instar larvae was very sensitive because the cuticle is not tanning and easy to germination tube of conida of fungus to penetrate integument to hymocel by using digestive enzymes and mechanical pressure on it Results showed that the mortality percentages of the larvae increased with an increased concentration of fungal suspension.

Assaf *et al.* (2011) when studying the fungus *Paecilouyces farinosus* on insect *Melasoma papule* the percentage of mortality to first instar larvae was 95% in concentration  $1 \times 10^8$  spore /ml. Jasim (2002) when studying fungus *B.bassiauq* on *Rhizopertha domonica* found that the efficacy increasing and also mortality when the concentration of fungus increase it was 96,94,84,16 % in concentration  $4 \times 10^5$ ,  $3 \times 10^5$ ,  $2 \times 10^5$  spore /ml. Elek (1993) say that the height mortality was in the first instar at larvae *Rhizopertha dominica* because it was sensitive to fungi including *B.bassiana*.

Table (2). Effect of fungus suspension of *B. bassiana* on first larval instar of *S. cerealella*

Treatment	%mortality Larvae after treatment	%Distorted larvae	%Forming pupae	% Emergence Adults
Control	3.0	0.0	85	84
$1 \times 10^2$	4.0	16.8	14.6	7.1
$1 \times 10^4$	9.6	18.2	6.6	0.0
$1 \times 10^6$	16.0	19.2	0.0	0.0
L.S.D	4.5	2.73	18.1	11.9

**Effect of fungus, suspension in three concentrations  $1 \times 10^2$ ,  $1 \times 10^4$ ,  $1 \times 10^6$  of *B. bassiana* on pupa of *S.ceralella* .**

Table (3) indicates that when was sprayed the suspension of *B.bassiana* the less percentage of emergence of adults was 8.3% in concentration  $1 \times 10^6$  spore /ml and the same percentage 8.3 % in concentration  $1 \times 10^4$  spore /ml while the heiest percentage of emergence in control treatment it was 96.3 % .

Albaita (2007) used suspension of fungus *B.bassiana* with concentration  $3.4 \times 10^{-6}$  spore /ml the mortality of pupa was 70% to insect *callsobruchus maculates* .Salih *et al.* (2010) used suspension of *B.bassiana* in concentration  $1 \times 10^{-1}$  spore /ml to control pupa of *musca domestica* and the mortality was ehibited 36.6 % .

Table (3) Effect of *B.bassiana* concentration  $1 \times 10^2$ ,  $1 \times 10^4$ ,  $1 \times 10^6$  spore /ml on pupa of *S.cerealella*

Treatment	% Formed pupae	% Emergence Adults
Control	96.3	75.2
$1 \times 10^2$	47.4	28.3
$1 \times 10^4$	35.5	8.3
$1 \times 10^6$	8.3	8.3
L.S.D 0.05	28.6	26.6

**Effect of *B.bassiana* concentrations  $1 \times 10^2$ ,  $1 \times 10^4$   $1 \times 10^6$  spore /ml on mortality adults *S. cerealella***

The data in table (4 ) the data in the was significant difference of mortality between three concentration of fungus  $1 \times 10^2$ ,  $1 \times 10^4$ ,  $1 \times 10^6$  spore /ml it were 43.3, 56.6, 70.0 % respectively and also difference found between concentration and control treatment.

After 8 hours the mortality of adults was 63, 73.3, 83.3 % respectively for three concentration respectively in control treatment was 3.3 % .

After 96 hours the mortality of adults treated with three concentration of fungus *B.bassiana* were 86.6, 86.6, 96.6 % respectively while in control was 16.6 % .

Table (4). Effect of *B.bassiana* concentration  $1 \times 10^2$ ,  $1 \times 10^4$ ,  $1 \times 10^6$  spore /ml on adults *S. cerealella*

Treatment	Adult mortality %		
	Exposure time(hour)		
	24	48	96
Control	3.3	3.3	16.6
$1 \times 10^2$	34.3	63.3	86.6
$1 \times 10^4$	56.6	73.3	86.6

$1 \times 10^6$	70.0	83.3	96.6
L.S.D 0.05	18.8	17.1	10.8

These differences in mortality among different instars may be related to enzymatic activity. It has been reported that the activity of detoxification enzymes varies considerably among and within developmental stages. The activity low in egg stage, increased with each larval instar and then declines to zero at pupation (Ahmad, was 1986; Mullin, 1988). Kaur et al (2011) mentiaed the virulence of *B. bassiana* was tested against second, third and 4th instar larvae of *S. litura* using three concentrations i.e.  $2.03 \times 10^8$ ,  $4.03 \times$

$10^6$  and these  $1.47 \times 10^5$  spores/ml, All treatments resulted in significantly higher mortality than control and ,higher mortality in second instar larvae than third and fourth instar larvae and gave negative effects of fungal infection of *Beauveria bassiana* on growth and development of *Spodoptera litura* resulted in significant reduction in adult emergence ,Likewise, Hafez et al. (1997) documented decrease in emergence of *P. operculella* , from 100 % in control to zero percent at  $16.5 \times 10^8$  conidia/ml

## References

- Al- yoni, A., 1993. Production and breeding of field crops – part one ministry of high education printing Baghdad university – college of Agriculture. 69, pp.148.
- Ahmad, S., 1986. Enzymatic adaptations of herbivorous insects and mites to phytochemicals. Journal of chemical ecology, 12(2), pp.533-560.
- Al-hydrai, A., 2002. Laboratories and fields studying on effect fungus *B . bassiana* on corn borer *Sesamia cretica* Master thesis , college of Agriculture, Baghdad university.
- Mohammad Ali, H.M. (2007). Study Effect of Ethanolic extract of leaf and frut of dorona *Durtarepens* and fungus *Beauveria bassiana* on biological aspect of mosquito *Culex pipien pipien* .Master thesis , college of science for woman , Baghdad university.
- Assaf, L. H., Feyroz, R. H., and Geban, H. Y., (2011). Evaluation of the Entomopathogenic fungi, *Beauveria bassiana* (bals) vuill and *Paecilomyces farinosus*(Dicks Exfr.) Against the poplar leaf Beetle *Melasoma populi*. Agricultural and veterinary science Duhok Univ. 14(1), pp.35-44.
- Bushra, S and Aslam, M., (2014). Management of *Sitotroga cerealella* (oliver) in stored *Beauveria bassiana* on development and reproductive potential of *Spodoptera chiffs*, Newjercoy.pp. 676.
- Carton, B. 1988. Development of genetically improved steains of *Bacillus thuringiensis*.cereal grains. Archives of phytopathology and pland protection. 47(19), pp.2365-2376.
- Development and utilization of cultivated plant. Pretice-Hall, Inc. Englewood development of *Rhyzopertha dominica* (F) (Coleoptera:Bostrychida) ,
- Elek, J.A.,1993. Effect of chitin- Synthesis-inhibitor, chlorfluazuron, on immature entomopathogenic fungus,*Beauveria bassiana* on the potato tuber moth
- Fornal, J., Jelinski, T., Sadowska, J., Grunda, S., Nawrot, J., Niewiada, A., litura (Fabricius ).*Journal of Biopesticides*, 4 (2), pp.161-168.
- Gindin, G., Ment, D., Rot, A., Glazer, I. and Samish, M., 2009. Pathogenicity of *Metarhizium anisopliae* (Hypocreales: Clavicipitaceae) to tick eggs and the effect of egg cuticular lipids on conidia development. *Journal of medical entomology*, 46(3), pp.531-538.
- Hafez, M., Zaki, F. N., Moursy, A. and Sabbour, M., (1997).Biological effects of How lander , A.J. and matin , A.S. 1988 Observations on the preharvest in formation of paddy by stored grain pests in Banglandesh . Journal

- of stored products Research 24 (4), pp. 229-231
- Hudson, T.H; William , I. F. and Anton, K. M., (1981). Plant science Growth, In :Hedin, P, Menu, J.and Hollingsworth, R. (Eds.), Biotechnology for crop
- Jasim , K.H., (2002) Effect of some Integrated biocontrol Factors to control lesser grain borer *Rhizopertha dominica* on rice seeds , *Journal of Agriculture science* (special issue) 7(5), pp.98-104.
- Kaur, S., Kaur, H.P., Kaur, K. and Kaur, A., 2011. Effect of different concentrations of *Beauveria bassiana* on development and reproductive potential of *Spodoptera litura* (Fabricius). *Journal of Biopesticides*, 4(2), pp.161-168.
- Lacey , L.A., 1997. Manual of techniques in insect pathology (Biological techniques) Academic press san diego . London .Boston. p.408 .
- Mullin, C. A., 1988. Adaptive relationships of epoxide hydrolase in herbivorous arthropods. *Journal of Chemical Ecology*,14. pp. 1867.
- Navon, A. and Ascher, K.R.S., 2000. Bioassays of Entomopathogenic Microbes and Nematodes CABI publishing pp322. *Phthorimaea operculella* (Seller). *Anz. Schadlingskde, Pflanzenschutz, phytochemicals. Journal of Chemical Ecology*, pp. 12: 533.
- Proceedings of the 6<sup>th</sup> International working. protection, American Chemical Society, Washington D. C., 260-279.
- Salih, H.A.M., Emad A.M. and Noor S. A., 2010. Effect of suspension of fungus *Beauveria bassiana* on live stage for house fly *Musca domestica* – *Iraqi Journal of Biotechnology*. 6(4),pp.690-700.
- SAS. (2012) Statistical Analysis System , User's Guide. Statistical. Version 9,1<sup>th</sup> ed SAS. Inst. Inc. Cary. N.USA.
- Waechalenski, J.R. and Blaszcak, W., 2007. Detection of granary weevil *Sitophilus granarius* L., eggs and internal stage analysis. *J.Stor.Prod.Res.*43:142-*Umweltschutz*, 70, pp.158-159.